

Next-to-leading order corrections to gauge-mediated gaugino masses

Marco Picariello and Alessandro Strumia

*Dipartimento di Fisica, Università di Pisa and
INFN, sezione di Pisa, I-56126 Pisa, Italia*

Abstract

We compute the next-to-leading order corrections to the gaugino masses M_i in gauge-mediated models for generic values of the messenger masses M and discuss the predictions of unified messenger models. If $M < 100$ TeV there can be up to 10% corrections to the leading order relations $M_i \propto \alpha_i$. If the messengers are heavier there are only few % corrections. We also study the messenger corrections to gauge coupling unification: as a result of cancellations dictated by supersymmetry, the predicted value of the strong coupling constant is typically only negligibly increased.

1 Introduction

The “*gauge mediation*” scenario for supersymmetric particle masses [1] can be realized in reasonable models [2, 3]. Furthermore, with a unified spectrum of messenger fields, it gives rise to some stable and acceptable prediction for the spectrum of supersymmetric particles. One of these predictions is the ‘unification prediction’ for the gaugino masses M_i ($i = 1, 2, 3$): at one-loop order the RGE-invariant ratio $\rho_i \equiv M_i/\alpha_i$ is the same for all the three factors of the SM gauge group $G_{\text{SM}} = \bigotimes_i G_i = \text{U}(1)_Y \otimes \text{SU}(2)_L \otimes \text{SU}(3)_C$. This prediction is sufficiently stable that it is interesting to compute it with more accuracy.

A detailed computation allows a comparison with the gaugino spectrum predicted by the alternative scenario known as “*unified supergravity*” [4]. Unification relations for the $\rho_i(E)$ are in fact the more stable prediction of this second scenario: gauge couplings and gaugino masses could receive sizeable GUT threshold corrections; but these corrections largely cancel out in the ratios $\rho_i(M_{\text{GUT}})$, since α_i and M_i have the same one-loop RGE evolution. The testable predictions for the low-energy running ρ_i ratios in the $\overline{\text{DR}}$ scheme [5] are (including NLO RGE corrections [6], but neglecting possible unknown $\mathcal{O}(\%)$ GUT-scale threshold effects [7])

$$\frac{\rho_1(M_Z)}{\rho_2(M_Z)} \approx 1.02, \quad \frac{\rho_2(M_Z)}{\rho_3(M_Z)} \approx 0.97. \quad (1)$$

This should be compared with the corresponding prediction in gauge-mediation models for the ratios ρ_i/ρ_j , plotted in fig.s 2 and 3.

The computation of gaugino masses with NLO precision is done in sections 2 and 3 for generic values of the messenger mass M . It requires the following main steps:

1. compute the renormalized running gaugino masses, $M_i(E_H)$ at $E_H \sim M$, in the effective theory without messengers.

2. compute the running gaugino masses, $M_i(E_L)$ at $E_L \sim M_Z$, evolving $M_i(E_H)$ down to the Fermi scale with 2 loop RGE equations.
3. use $M_i(E_L)$ to compute directly measurable quantities, like the gaugino pole masses, M_i^{pole} , including all 1 loop effects at the electroweak scale.

The computation necessary for step 1 is done in section 2. We will employ supersymmetric dimensional regularization, so that the renormalization scale E will be the $\overline{\text{DR}}$ scale, $\bar{\mu}$. The RGE necessary for step 2 (recalled in appendix B) can be read from the literature [6]. The one-loop expressions for pole gaugino masses in terms of running parameters are also well known [8]. Since various unmeasured and unpredicted parameters (like the so-called μ -term) would enter the final step 3, we prefer to show our final predictions for the running MSSM gaugino masses renormalized at $\bar{\mu}_L = M_Z$, without including the gauge corrections at the electroweak scale.

We will compute these predictions in *unified messenger models*. One more step is necessary to impose the unification constraints on the messenger spectrum, namely

0. compute the messenger spectrum, $M_n(E_H)$, evolving the unified $M_n(M_{\text{GUT}})$ down to the messenger scale $E_H \sim M_n$ with 2 loop RGE equations.

The necessary RGE equations are given in appendix B. In sec. 3 we study the predictions of gauge mediation models with an unified messenger spectrum. If the messenger spectrum is only negligibly splitted by supersymmetry breaking effects, the NLO corrections to the LO unification-like relations $M_i \propto \alpha_i$ are around few % and numerically not much different from the ones present in unified supergravity models. Larger effects (up to 10%) can be present if the messengers are very light, $M \lesssim 50 \text{ TeV}$.

We also study the corrections to gauge coupling unification due to the presence of messenger fields below the unification scale. Messenger threshold effects largely cancel messenger corrections to two loop RGE running, as dictated by supersymmetry [9, 10].

2 Computation

In this section we do the computations necessary for step 1. We assume that the messenger fields sit in real representations $R = \bigoplus_n R_n$ of the SM gauge group and have a supersymmetric mass term M_n together with a non-supersymmetric mass term F_n . For the moment we assume that the messengers lie in self-conjugate complex representations, $R_n = X_n \oplus \bar{X}_n$. See appendix A for a more detailed discussions of the notations, of the model, of its spectrum, and of the relevant Lagrangian.

In order to compute the running gaugino masses $M_i(\bar{\mu})$ at $\bar{\mu} \lesssim M_n$ in the effective theory we first compute the gauge-independent *pole* gaugino masses in the two versions of the theory. The computation in the full theory (with messengers) is done in section 2.1 — the computation in the effective theory (with messengers integrated out) is done in section 2.2. Requiring that the two theories describe the same gaugino masses up to second order in α_i we get, in section 2.3 the gaugino mass terms in the effective theory.

We employ the Feynman-Wess-Zumino gauge and the supersymmetric $\overline{\text{DR}}$ regularization [5] in both versions of the theory. The final $\overline{\text{DR}}$ values of the gauge couplings α_i and of the gaugino masses M_i can be converted into the $\overline{\text{MS}}$ ones (i.e. the ones obtained with naïve dimensional regularization) using

$$\alpha_{\overline{\text{MS}}} = \alpha_{\overline{\text{DR}}} \left(1 - \frac{C(G)}{3} \frac{\alpha}{4\pi} \right), \quad M_{\overline{\text{MS}}} = M_{\overline{\text{DR}}} \left(1 + C(G) \frac{\alpha}{4\pi} \right)$$

where the group factors are defined as follows. For each representation R of a gauge group $G = \bigotimes_i G_i$ we define the “Dinkin index” $T_i(R)$ and the “quadratic Casimir” $C_i(R)$ in terms of the generators T_{Ri}^a as

$$\sum_a T_{Ri}^a \cdot T_{Ri}^a = C_i(R) \mathbb{I}, \quad \text{Tr } T_{Ri}^a T_{Rj}^b = T_i(R) \delta_{ij} \delta^{ab}. \quad (2)$$

With generators canonically normalized so that $T(\mathbf{n}) = T(\bar{\mathbf{n}}) = 1/2$ for the fundamental $\mathbf{n}$ representation of a $\text{SU}(n)$ group, the quadratic Casimir of the adjoint representation of a $\text{SU}(n)$ group is $C(G) = T(G) = n$, while

graph Γ and its value V^Γ		graph Γ and its value V^Γ	
XX	$\alpha_j C_j(X)(-\frac{10}{3})$	$X\tilde{X}$	$\alpha_j [C_j(X) - \frac{1}{2}\delta_{ij}C_i(G)](2/\varepsilon + 4)$
$\widetilde{XX}$	$\alpha_j C_j(X)\frac{4}{3}(-1/\varepsilon + 1)$	$\widetilde{X\tilde{X}}$	$\alpha_j [C_j(X) - \frac{1}{2}\delta_{ij}C_i(G)](-4)$
$\tilde{X}\tilde{X}$	$\alpha_j C_j(X)(3/\varepsilon - 2/3)$	$X\lambda$	$\alpha_i C_i(G)(4/\varepsilon - 1)$
$\widetilde{\tilde{X}\tilde{X}}$	$\alpha_j C_j(X)\frac{8}{3}(-2/\varepsilon - 1)$	$\tilde{X}\lambda$	$\alpha_i C_i(G)(1/\varepsilon - 1)$
$\tilde{X}\tilde{X}'$	$\alpha_j C_j(X)(\frac{5}{3}\frac{1}{\varepsilon} + \frac{4}{3})$	$\lambda\lambda$	$\alpha_i C_i(G)(-4/\varepsilon_{\text{ir}} + 4)$

Table 1: Contributions $\delta M_i^\Gamma = M_i^{(1)} V^\Gamma / 4\pi$ of the single graphs Γ in the limit $F \ll M^2$.

$C(G) = 0$ for G a U(1) factor. The values of the coefficients for the SM gauge group and for representations contained in $5 \oplus \bar{5}$ and $10 \oplus \bar{10}$ representation of SU(5) are given in table 2.

2.1 Computation in the full theory

The NLO correction to the pole gaugino masses are given by the ten two-loop diagrams shown in figure 1 and some one-loop renormalization factor. It is convenient to separate the renormalization factors due to the light MSSM loops from the ones due to messenger loops. We write the various contributions to the pole gaugino masses as

$$M_i|_{\text{pole}}^{\text{full}} = \hat{M}_i^{(1)} + \sum_{\Gamma} \delta M_i^\Gamma, \quad \hat{M}_i^{(1)} \equiv \sum_n \hat{M}_{in}^{(1)} = \frac{\alpha_i}{4\pi} \sum_n \frac{F_n}{M_n} T_i(R_n) \hat{g}_1(x_n) = M_i^{(1)} + \mathcal{O}(\varepsilon) \quad (3)$$

where $x_n = F_n/M_n^2$ and the sum $\sum_n$ extends over all the messengers. The one-loop function $\hat{g}_1(x) \equiv g_1(x) + \varepsilon \delta g_1(x) + \mathcal{O}(\varepsilon^2)$ is [11]

$$\hat{g}_1(x) = \frac{1+x}{x^2} \ln(1+x) \left\{ 1 + \varepsilon \left[1 - \frac{\ln(1+x)}{2} \right] \right\} (1 + \varepsilon \ln \frac{\bar{\mu}^2}{M^2}) + (x \rightarrow -x) + \mathcal{O}(\varepsilon^2) = 1 + \frac{x^2}{6} + \mathcal{O}(x^4, \varepsilon) \quad (4)$$

All parameters are unrenormalized ('bare'). Here we list all the contributions to the pole gaugino masses.

- The contribution given by the sum of the **two loop diagrams** of fig. 1. They give

$$\begin{aligned} \delta M_i^{2 \text{ loop}} &= \frac{\alpha_i}{4\pi} \sum_n \frac{F_n}{M_n} T_i(R_n) \left\{ \frac{\alpha_i}{4\pi} C_i(G) \left[\hat{g}_1(x_n) \left(\frac{4}{\varepsilon_{\text{uv}}} - \frac{4}{\varepsilon_{\text{ir}}} + g_{\text{IR}} \right) + 2g_2(x_n) \right] + \right. \\ &\quad \left. + \sum_{j=1}^3 \frac{\alpha_j}{4\pi} C_j(X_n) \left[g_C(x_n) - 4g_1(x_n) - \frac{2x_n \hat{g}_1'(x_n)}{\varepsilon_{\text{uv}}} \right] \right\} \left(\frac{\bar{\mu}^2}{M_n^2} \right)^\varepsilon \end{aligned} \quad (5)$$

where $\hat{g}_1'(x)$ is the derivative of $\hat{g}_1(x)$ with respect to x , and the functions g_2 and g_C are

$$\begin{aligned} g_2(x) &= \frac{\ln(1+x)}{2x^2} [2(1+x) + \ln(1-x) + (2x+2+1/x)\ln(1+x)] + \\ &\quad + \frac{1}{x} [\text{Li}_2(\frac{2x}{1+x}) - 2\text{Li}_2(x)] + (x \rightarrow -x) \end{aligned} \quad (6a)$$

$$g_C(x) = \frac{\ln(1+x)}{x^2} [8 + 6x + \ln(1-x) + (1-x+2/x)\ln(1+x)] + (x \rightarrow -x) \quad (6b)$$

In the limit $F \ll M^2$ ($x \rightarrow 0$) $g_1(0) = g_2(0) = 1$ and $g_C(0) = 0$. We have denoted as $1/\varepsilon_{\text{uv}}$ an $1/\varepsilon$ ultraviolet (UV) pole, and as $1/\varepsilon_{\text{ir}}$ a pole of infrared (IR) origin. In all the graphs it is possible to set the external gaugino momentum to zero, except in the infrared divergent $\lambda\lambda$ graph of fig. 1. It is convenient to split it into a part computed with zero external momentum, that contributes to g_2 , plus the remainder, that gives the g_{IR} term in eq. (5). The value of g_{IR} coincides with the 'non naïve part' of the 'asymptotic




Figure 1: The diagrams that contribute to the NLO gauge corrections to gauge mediated gaugino masses. The thick continuous (dashed) lines represent the fermionic (bosonic) messengers. The thin wavy (continuous) lines represent the gauge bosons (gauginos).

expansion¹ in the external gaugino momentum, and can be seen as the contribution of the $\lambda\lambda$ graph with the heavy messenger loop contracted to a point. This technical detail is useful, because a corresponding one-loop diagram gives the same contribution to the effective theory, so that we do not need to compute g_{IR} .

- **On-shell renormalization of the gaugino wave function:** the renormalized gaugino field is $\lambda_i = (1 + z_i + z'_i)^{1/2} \cdot \lambda_i|_{\text{bare}}$ with

$$z'_i = \frac{\alpha_i}{4\pi} \sum_n T_i(R_n) \left\{ \frac{1}{\varepsilon} + \ln \frac{\bar{\mu}^2}{M_n^2} + \frac{x_n^2 + (1 - x_n^2)[\ln(1 - x_n) + \ln(1 + x_n)]}{2x_n^2} \right\}.$$

We have separated this correction into two parts: z due to all the MSSM particles, and z' due to messenger loops only. The corresponding corrections to the gaugino masses, $\delta M_i^z + \delta M_i^{z'}$, are obtained expressing

¹ The general technique of asymptotic expansions of Feynman diagrams is described in [12]; a much simpler discussion, sufficient for the purposes of this computation, can be found in [13], where an accurate distinction between UV and IR divergences is made.

rep.	C_1	C_2	C_3	T_1	T_2	T_3
$d \oplus \bar{d}$	2/15	0	8/3	2/5	0	1
$L \oplus \bar{L}$	3/10	3/2	0	3/5	1	0
$Q \oplus \bar{Q}$	1/30	3/2	8/3	1/5	3	2
$u \oplus \bar{u}$	8/15	0	8/3	8/5	0	1
$e \oplus \bar{e}$	6/5	0	0	6/5	0	0

Table 2: Values of the group factors for the G_{SM} fragments of the $5 \oplus \bar{5}$ and $10 \oplus \bar{10}$ $\text{SU}(5)$ representations.

the LO result in terms of the renormalized field λ_i : $\delta M_i^{z'} = -\hat{M}_i^{(1)} \cdot z'_i$. We do not need to specify the MSSM part because it is the same in both versions of the theory.

- **Renormalization of the gauge couplings:** we choose to express the bare gauge couplings of the *full* theory, $g_i|_{\text{bare}}$, as function of the quantum-corrected gauge couplings in the *effective* theory renormalized in the $\overline{\text{DR}}$ scheme, $g_i(\bar{\mu})$. Defining $\alpha_i \equiv g_i^2/4\pi$ and including the messenger thresholds effects we find

$$\alpha_i(\bar{\mu}) = \alpha_i|_{\text{bare}} - \delta\alpha_i - \delta'\alpha_i, \quad \frac{\delta'\alpha_i}{\alpha_i} = \frac{\alpha_i}{4\pi} \sum_n T_i(R_n) \left\{ \frac{1}{\varepsilon} + \ln \frac{\bar{\mu}^2}{M_n^2} - \frac{\ln(1-x_n) + \ln(1+x_n)}{6} \right\}.$$

We have separated the correction to α_i into two parts: $\delta\alpha$ due to all the MSSM particles, and $\delta'\alpha$ due to messenger loops only. The corresponding corrections to the gaugino masses, $\delta M_i^{\delta\alpha} + \delta M_i^{\delta'\alpha}$, are obtained expressing the LO result in terms of the renormalized MSSM gauge coupling $\alpha_i(\bar{\mu})$: $\delta M_i^{\delta'\alpha} = \hat{M}_i^{(1)} \cdot \delta'\alpha_i/\alpha_i$. We do not need to specify the MSSM part because it is the same in both versions of the theory. Notice that the messenger contribution $\delta M_i^{\delta'\alpha} + \delta M_i^{z'}$ is UV finite.

- **Renormalization of F and M .** We employ $\overline{\text{DR}}$ renormalization: F and M are defined as their bare values plus the pole parts of their quantum corrections. As a consequence of the non-renormalization theorem, the parameters M_n and F_n renormalize in the same way:

$$\{F_n, M_n\} = \left(1 - \sum_j \frac{\alpha_j}{4\pi} 2C_j(X_n)\right) \{F_n, M_n\}_{\text{bare}}$$

The corrections to the gaugino masses are obtained expressing the bare parameters in terms of the renormalized ones in the one loop result:

$$\delta M_i^{F,M} = \frac{\alpha_i}{4\pi} \sum_n \frac{F_n}{M_n} T_i(R_n) \sum_j \frac{\alpha_j}{4\pi} C_j(X_n) \left\{ 4g_1(x_n) + \frac{2x_n \hat{g}'_1(x_n)}{\varepsilon_{\text{uv}}} \right\}. \quad (7)$$

The first term of (7) derives from $\ln \bar{\mu}^2/M^2$ in (4) and, for $x = 0$, cancels the NLO correction to M_i proportional to α_j produced by the two-loop diagrams.

As an aside remark, it could be of interest to know that the NLO squared pole mass of the lightest scalar messenger, M_{n-}^2 , is

$$\begin{aligned} \frac{M_{n-}^2}{M_n^2 - F_n} &= 1 - \sum_j \frac{\alpha_j}{4\pi} C_j(X_n) \left\{ 2(x_n - 2)(2 + \ln \frac{\bar{\mu}^2}{M_n^2}) + \right. \\ &\quad \left. + \frac{2x_n - 3 - 3x_n^2}{x_n - 1} \ln(1 - x_n) + \frac{2x_n^2}{x_n - 1} \ln x_n - (1 + x_n) \ln(1 + x_n) \right\} \end{aligned}$$

when expressed in terms of $\overline{\text{DR}}$ parameters.

2.2 Computation in the effective theory

In the effective theory we have to compute the pole gaugino masses in terms of the coefficients of the running gaugino mass term operator, $-M_i(\lambda_i\lambda_i + \text{h.c.})$, expanded as a series in the gauge couplings:

$$M_i = \frac{\alpha_i}{4\pi} M_i^{(1)} + \frac{\alpha_i}{4\pi} \frac{\alpha_j}{4\pi} M_{ij}^{(2)} + \dots \quad (8)$$

where $M_i^{(1)}$ are the known LO coefficients, and $M_{ij}^{(2)}$ are the NLO coefficients that we want ultimately to extract. The pole gaugino masses at $\mathcal{O}(\alpha^2)$ order are given by

- the contribution from the renormalization of the gaugino wave function. This coincides with the δM_i^z correction present also in the full theory.
- the contribution from the renormalization of the gauge couplings. This coincides with the $\delta M_i^{\delta\alpha}$ correction present also in the full theory.
- a one loop diagram (gauge correction to the gaugino propagator). As said, it is not difficult to see that it gives the same contribution of the asymptotic expansion in the external gaugino momentum of the two-loop $\lambda\lambda$ diagram of fig. 1.

More in detail the pole gaugino masses in the effective theory are

$$M_i^{\text{pole}} = \frac{\alpha_i}{4\pi} M_i^{(1)} \left\{ 1 + \frac{\alpha_i}{4\pi} C_i(G) \left(\frac{4}{\varepsilon_{\text{uv}}} - \frac{4}{\varepsilon_{\text{ir}}} + g_{\text{IR}} \right) \right\} + \frac{\alpha_i}{4\pi} \frac{\alpha_j}{4\pi} M_{ij}^{(2)} + \delta M_i^z + \delta M_i^{\delta\alpha}. \quad (9)$$

A further simplification occurs. The sum of the three effective-theory quantum corrections, all proportional to $M_i^{(1)}$, is both infrared and ultraviolet convergent (because the combination M_i/α_i is RGE-invariant at one loop). For this reason we do not need to worry about the $\mathcal{O}(\varepsilon)$ terms that distinguish $\hat{M}_i^{(1)}$ from $M_i^{(1)}$.

2.3 Matching

The matching procedure is particularly simple: the running gaugino masses in the effective theory at NLO order are simply given by the full theory result, omitting those quantum corrections that are present also in the effective theory result, eq. (9). The MSSM running gaugino masses at NLO order are

$$M_i = \frac{\alpha_i}{4\pi} \sum_n \frac{F_n}{M_n} T_i(R_n) \left\{ g_1(x_n) + \frac{\alpha_i}{4\pi} \left[2C_i(G)g_2(x_n) + \sum_m T_i(R_m)g_T(x_m) \right] + \sum_{j=1}^3 \frac{\alpha_j}{4\pi} C_j(X_n) \left[g_C(x_n) - 2x_n g_1'(x_n) \ln \frac{\bar{\mu}^2}{M_n^2} \right] + \frac{\lambda_n^2}{(4\pi)^2} g_\lambda(x_n) \right\}. \quad (10)$$

The parameters are renormalized as discussed in sec. 2.1. The last term is the effect of a possible Yukawa coupling $\lambda_n S X_n \bar{X}_n$ in the superpotential, where S is a gauge singlet². The functions g_1 , g_2 and g_C have been defined in eq.s (4) and (6), while g_T and g_λ are

$$g_T(x) = \frac{2x^2 - 3}{6x^2} \ln(1 - x^2) - \frac{1}{2}, \quad (11a)$$

$$g_\lambda(x) = -\frac{\ln(1+x)}{2x^3} [2x(3+2x) - x \ln(1-x) + (5+3x) \ln(1+x)] + \frac{x-4}{x^2} \text{Li}_2(x) + (x \rightarrow -x). \quad (11b)$$

The functions g_1 and g_2 are normalized such that $g_1(0) = g_2(0) = 1$, while $g_C(0) = g_T(0) = g_\lambda(0) = 0$. So far we have assumed that $R_n = X_n \oplus \bar{X}_n$. If there are also messenger fields Σ in a real representation R_Σ , the appropriate group factors are $T(R_\Sigma) = T(\Sigma)$ and $C(X) \rightarrow C(\Sigma)$.

²The computation of Yukawa corrections, that we do not present, involves one new feature: a renormalization of F_n/M_n . For simplicity, we have given the Yukawa correction assuming that all the F_n and M_n are produced by a vacuum expectation value of the superfield S . In this case the new feature becomes an irrelevant common renormalization of F_n/M_n . The corresponding effect in the final result, eq. (10), has been absorbed in an appropriate (non $\overline{\text{DR}}$) renormalization of F_n .




Figure 2: *NLO predictions for the measurable ratios $\rho_i(M_Z)/\rho_j(M_Z)$ ($\rho_i \equiv M_i/\alpha_i$) in the minimal gauge mediated model ($n_5 = 1$, $n_{10} = 0$) for generic values of the unified messenger mass. For comparison, we also show the prediction of unified supergravity models (neglecting possible small GUT-scale effects).*

In the limit $x_n \rightarrow 0$ the NLO prediction for the running gaugino masses does not depend on the messenger spectrum

$$M_i(\bar{\mu}) \stackrel{F \ll M^2}{\approx} \frac{\alpha_i(\bar{\mu})}{4\pi} \sum_n \frac{F_n}{M_n} T_i(R_n) \left[1 + \frac{\alpha_i}{4\pi} 2C_i(G) \right] \quad \text{at } \bar{\mu} \sim M_n. \quad (12)$$

We remember that $C_i(G) = \{0, 2, 3\}$. This prediction for the three gaugino masses, without knowing the values of the F_n/M_n parameters, is of interest only if their number is less than three. This happens, for example, if the messengers lie in a $5 \oplus \bar{5}$ representation of $SU(5)$.

3 Predictions of unified messenger models

We will show the NLO predictions in models where the messenger spectrum satisfies unification relations. These models are not only more predictive but also more appealing: the successful unification of the gauge couplings is not destroyed and a unified messenger spectrum helps in avoiding undesired one-loop contributions to sfermion masses. To be more specific we assume that the messengers fill n_5 copies of $5 \oplus \bar{5}$ and n_{10} copies of $10 \oplus \bar{10}$ representations of the unified group $SU(5)$, so that the messenger contribution to the one loop coefficient of the gauge β functions is $T_i(R) = n_5 + 3n_{10}$. We assume that the messenger mass parameters M_n and F_n arise from the vacuum expectation value of one $SU(5)$ -singlet field S coupled to the messengers via Yukawa interactions $\lambda_n S X_n \bar{X}_n$. Imposing the unification relations the running $\overline{DR}$ mass parameters at $\bar{\mu} \sim M_n$ are thus obtained with NLO precision via two-loop RGE evolution from M_{GUT} down to $\bar{\mu}$:

$$x_n(\bar{\mu}) = \frac{F_n}{M_n^2} = \mathcal{U}_n(M_{GUT} \rightarrow \bar{\mu})[x_N(M_{GUT}) + \delta_n x_N] \quad \text{for messengers } R_n \text{ unified in } R_N$$

where $\delta_n x_N$ represent unknown one-loop threshold effects at the unified scale, that we will neglect. ‘Reasonable’ threshold effects give small corrections also when $x \sim 1$. We also neglect NLO Yukawa corrections, possibly




Figure 3: As in fig. 2, but with a ‘less minimal’ messenger content.

relevant when $x \sim 1$. The NLO RGE equations for M_n and F_n are given in appendix B (the combination F_n/M_n is RGE-invariant, and is not corrected by threshold effects). Already at LO, the RGE evolution of the messenger spectrum gives corrections of relative order $\mathcal{O}(x^2 \alpha \ln M_{\text{GUT}}/M)$ to the relations $M_i \propto \alpha_i$: for light messengers the leptonic messengers are approximately 2 times lighter than the hadronic ones. This explains the larger effect present for light messengers ($x_N \sim 1$)³.

Fig. 2 shows the prediction of the minimal model with $(n_5, n_{10}) = (1, 0)$ for the measurable ratios ρ_i/ρ_j (the experimental errors on the gauge couplings negligibly affect the predictions for the $\rho_i(M_Z) \equiv M_i(M_Z)/\alpha_i(M_Z)$). In this figure we have considered the whole range of possible messenger masses, distinguishing the smaller values of the messenger mass for which the lightest supersymmetric particle (LSP) decays before escaping detection, from the higher values for which the LSP decay is so slow (in absence of R -parity breaking) that destroys the nucleosynthesis products [3].

In fig. 3 we consider models with more than a single unified messenger. For the sake of illustration we have combined their contributions assuming that all the messengers have the same unified M_n and F_n . For values of M higher than the ones considered in fig. 3 ($x \ll 1$) all neglected NLO terms are completely irrelevant, the prediction does not depend on the messenger content, and is the same as in fig. 2. In all plots we have fixed the F -terms requiring that the running gluino mass be $M_3(M_Z) = 500$ GeV. Any other reasonable value of the gluino mass gives the same prediction for ρ_i . We remember that we have *not* included the one-loop corrections at the electroweak scale to M_i , that depend on unmeasured (but measurable) and unpredicted parameters (mainly the μ -term). The error on these predictions, due to remaining NNLO effects, is estimated to be at the *per-mille* level, much smaller than the expected experimental error on the gaugino masses.

In fig.s 2, 3 we have also plotted the corresponding NLO prediction of unified supergravity models, without including unknown possible GUT-scale corrections. The unification relation $\rho_i \propto \mathbb{I}$ is in fact only corrected at the % level by GUT-scale threshold [7] and gravitational [15] effects, that could instead give much larger corrections to the unification relations for the α_i and for the M_i . The RGE contribution from the top A term,

³ Values of $x_N > 1$ (negative squared tree-level masses for scalar messengers) give an acceptable spectrum of physical messenger masses ($x_n(\bar{\mu}) < 1$), leaving only charge and/or colour breaking (CCB) minima at very large field values. This situation is not forbidden. On the contrary, non-dangerous CCB minima are quite generic in gauge mediated models [14].




Figure 4: *Percentage correction to the unification prediction for $\alpha_3(M_Z)$ due to the presence of messengers unified in the simplest representations of SU(5).*

driven towards its IR fixed point value, $A_t(M_Z) \approx 2M_2$, is also numerically negligible [6]. The same can be said for the bottom and τ contributions, that remain negligible also if $\tan\beta$ is large.

At this point it is also interesting to discuss the NLO correction that the presence of messenger fields gives to the *unification prediction for $\alpha_3(M_Z)$* . The percentage correction to $\alpha_3(M_Z)$ is plotted in fig. 4 for models with different messenger content, assuming that all different messengers have a common unified value of the M and F parameters (if $F \ll M^2$ it is only necessary to assume that the various M_n have the same order of magnitude). If $n_5 + 3n_{10} > 5$ too light messengers give a non-perturbative value of the unified gauge coupling.

It is interesting to see more in detail why this correction is much smaller than its naïve expectation and why it exhibits some curious property. For given values of the unification scale and of the unified gauge coupling, the low energy gauge couplings receive two contributions due to the presence of messengers: $\delta\alpha_i^{-1}(M_Z) = \delta_i^{\text{th}} + \delta_i^{\text{RGE}}$. The first contribution, δ_i^{th} , is due to messenger thresholds (gauge corrections distort the unified messenger spectrum); the second one is due to messenger corrections to the running of the gauge couplings (we can reabsorb the one-loop contribution in the definition of the unified gauge coupling, and consider only the two-loop contribution). In a sufficiently accurate approximation the two corrections are

$$\delta_i^{\text{RGE}} = \frac{1}{4\pi} \sum_j \left\{ b_{ij}^{(2)} \ln r_j - b_{ij}^{(2)\text{MSSM}} \ln r_j^{\text{MSSM}} \right\} \quad (13a)$$

$$\delta_i^{\text{th}} = \frac{1}{4\pi} \sum_n \sum_{p \in R_n} b_i^p \ln \frac{\bar{\mu}^2}{M_p^2} \stackrel{F \ll M^2}{=} -\frac{1}{4\pi} \sum_j \sum_n 4T_i(R_n) C_j(X_n) \ln r_j \quad (13b)$$

where $\sum_p$ extends over all the fermionic and scalar messengers, b_i^p is the contribution of a given messenger to $b_i^{(1)} = b_i^{\text{MSSM}} + \sum_p b_i^p$, and $b_i^{(1)}$ and $b_{ij}^{(2)}$ are the one and two-loop coefficients of the gauge β -functions in presence of messengers (explicitly given in appendix B). We define $r_i \equiv (1 - b_i \alpha_i(\bar{\mu}) / (4\pi) \ln M_{\text{GUT}}^2 / \bar{\mu}^2)^{-1/b_i} > 1$ while r_i^{MSSM} is its value without messengers. The overall correction to the unification prediction for the strong coupling constant is

$$\delta^{\text{mess}} \alpha_3(M_Z) = \delta^{\text{th}} \alpha_3 + \delta^{\text{RGE}} \alpha_3 = -\alpha_3^2(M_Z) (\delta_i^{\text{th}} + \delta_i^{\text{RGE}}) \Pi_i \quad (14)$$

where $\Pi_i = \{5/7, -12/7, 1\}$. The two corrections, $\delta^{\text{th}}\alpha_3$ and $\delta^{\text{RGE}}\alpha_3$ can be quite large ($\pm\mathcal{O}(0 \div 20)\%$) and depend separately on n_5 and on n_{10} . However the sum of the two contributions, plotted in fig. 4, is *much smaller*, typically positive, $\delta\alpha_3 = (0 \div 3)\%$, and depends only on the correction to the one-loop β -function coefficient $b_i^{\text{mess}} = n_5 + 3n_{10}$ (this is not true in the limit $x \sim 1$, where supersymmetry-breaking effects become relevant). This cancellation can be seen summing the expression for δ_i^{th} in the limit $F \ll M$, eq. (13b) (in which we have inserted the messenger masses M_n obtained via one-loop RGE evolution), with the RGE correction (in which we insert the general values of the two-loop β -function coefficients, written in eq. (B.2)):

$$\delta_i^{\text{mess}} = \delta_i^{\text{th}} + \delta_i^{\text{RGE}} \stackrel{F \ll M^2}{=} \frac{1}{4\pi} \left\{ 2C(G_i)b_i^{\text{mess}} \ln r_i + b_{ij}^{(2)\text{MSSM}} \ln \frac{r_j}{r_j^{\text{MSSM}}} \right\}. \quad (15)$$

Corrections due to possible messenger Yukawa couplings would cancel out. These cancellations reproduce the exact result found by J. Hisano and M. Shifman in [9, 10] working in toy models with the holomorphic supersymmetric gauge couplings. As shown in [16] this same reason is at the basis of the analogous cancellation between RGE and threshold effects encountered in our NLO computation of gaugino masses at $F \ll M$.

4 Conclusion

We have computed the next-to-leading order corrections to gaugino masses in gauge-mediated models for generic values of the messenger masses M . In unified messenger models there are up to 10% corrections to the unification-like relations $M_i(\bar{\mu}) \propto \alpha_i(\bar{\mu})$ between the running gaugino masses and the gauge couplings, but only if the messengers are strongly splitted by supersymmetry breaking. If instead $M > 100$ TeV there are only small (few %) corrections to the leading-order approximation $M_i \propto \alpha_i$, as shown in figs 2 and 3.

We have also studied the messenger corrections to gauge coupling unification. As a result of cancellations, dictated by supersymmetry, between large RGE and threshold corrections the predicted value of the strong coupling constant is typically only negligibly increased, as shown in fig 4.

In the limit $M^2 \gg F$ (heavy messengers) the same NLO prediction for gaugino masses, together with NLO predictions for sfermion masses, can be obtained [16] combining the techniques described in [17] and [9, 10]. If $F \sim M^2$ it is more difficult to obtain a NLO prediction of sfermion masses; however the LO results [11] show that the effects of large supersymmetry breaking in the messenger spectrum are much less relevant in the sfermion sector than in the gaugino sector.

Acknowledgements We are grateful to Riccardo Barbieri for many discussions, and to Riccardo Rattazzi for having pointed out a missing term (!) in the renormalization of our result.

A Relevant Lagrangian in quadri-spinor notation

We consider a theory with messenger chiral superfields Φ_L and Φ_R in self-conjugate complex representations of the SM gauge group (with generators T and $-T^T$). In presence of the superpotential

$$W = \int d^2\theta \lambda S \Phi_L \Phi_R, \quad \lambda \langle \hat{S} \rangle = M + \theta\theta F.$$

The messenger superfields $\Phi = A + \sqrt{2}\theta\psi + \dots$ contain the following mass eigenstates: messenger fermions ψ_L, ψ_R with a Dirac mass M , and pseudoscalar (scalar) messengers $A_{\pm} \equiv (A_L \pm A_R^*)/\sqrt{2}$ with mass $M_{\pm}^2 = M^2 \pm F \equiv M^2(1 \pm x)$. The supersymmetric gaugino Lagrangian is

$$\mathcal{L}_{\lambda} = \bar{\lambda}\bar{\sigma}^{\mu}iD_{\mu}\lambda - \frac{m}{2}(\lambda\lambda + \bar{\lambda}\bar{\lambda}) - \sqrt{2}g(A_L^{\dagger}T\psi_L\lambda + \bar{\lambda}\bar{\psi}_LT A_L - \lambda\psi_R^T T A_R^* - A_R^T T \bar{\psi}_R^T \bar{\lambda})$$

where D_{μ} is the standard gauge-covariant derivative and λ, ψ_L and ψ_R are Weyl fermions with the same chirality.

We want to rewrite the Lagrangian in terms of mass eigenstates. In order to employ our `Mathematica` [18] code for analytic computation of Feynman graphs, we need to write the messenger fermions as Dirac quadri-spinors Ψ and the gauginos λ as Majorana spinors Λ :

$$\Psi \equiv \begin{pmatrix} \psi_L \\ \bar{\psi}_R^T \end{pmatrix}, \quad \Lambda = \begin{pmatrix} \lambda \\ \bar{\lambda} \end{pmatrix} = \Lambda^c \equiv C\bar{\Lambda}^T$$

where C is the charge-conjugation matrix. The gaugino Lagrangian becomes

$$\mathcal{L}_\lambda = \frac{1}{2}\bar{\Lambda}(i\not{D} - m)\Lambda - g(A_-^\dagger T\bar{\Lambda}\Psi + \bar{\Psi}\Lambda T A_- - A_+^\dagger \bar{\Lambda}\gamma_5\Psi + \bar{\Psi}\gamma_5\Lambda T A_+) \quad (\text{A.1})$$

and the D terms become

$$D^a = A_L^* T^a A_L - A_R^* T^a A_R + \dots = A_+^* T^a A_- + A_-^* T^a A_+ + \dots$$

The gaugino $\Lambda\bar{\Lambda}$ propagator is a standard fermion propagator. It is possible to show that graphs with $\Lambda\Lambda$, $\bar{\Lambda}\bar{\Lambda}$ and $\bar{\Lambda}\Lambda$ propagators have the same value of the standard ‘ $\Lambda\bar{\Lambda}$ ’, by appropriately rewriting the vertices in terms of charge-conjugated fields⁴. This shows that the gaugino can be treated as an ordinary fermion field, but with symmetry factors computed like the ones of a *real* field.

In a general theory there will be several messenger pairs, each one with its M_n , F_n and $x_n \equiv F_n/M_n^2$.

B RGE evolution

The necessary RGE can be read from the literature [6]. The RGE equations for gauge couplings and gaugino masses in the $\overline{\text{DR}}$ scheme are

$$\begin{aligned} \frac{d}{dt} \frac{1}{g_i^2} &= b_i^{(1)} + \frac{1}{(4\pi)^2} \left[\sum_j b_{ij}^{(2)} g_j^2 - \sum_a b_{ia}^{(2)} \lambda_a^2 \right] \\ \frac{d}{dt} M_i &= -g_i^2 b_i^{(1)} M_i + \frac{g_i^2}{(4\pi)^2} \left[-\sum_j b_{ij}^{(2)} g_j^2 (M_i + M_j) + \sum_a b_{ia}^{(2)} \lambda_a^2 (M_i - A_a) \right] \end{aligned}$$

where $t(E) \equiv (4\pi)^{-2} \ln M_{\text{GUT}}^2/E^2$ and a runs over the third generation particles, $a = \{t, b, \tau\}$. In a general supersymmetric model with $\{\Phi\}$ matter superfields the coefficients are

$$b_i^{(1)} = -3C(G_i) + \sum_\Phi T_i(\Phi), \quad b_{ij}^{(2)} = 2C(G_i)b_i^{(1)}\delta_{ij} + 4\sum_\Phi T_i(\Phi)C_j(\Phi). \quad (\text{B.2})$$

In the models under consideration the field content is given by the MSSM fields plus n_5 copies of $5 \oplus \bar{5}$ and n_{10} copies of $10 \oplus \bar{10}$ $\text{SU}(5)$ messenger multiplets (in the effective theory without messengers, the coefficients are obtained taking $n_5 = n_{10} = 0$). The values of the coefficients are

$$\begin{aligned} b_i^{(1)} &= \begin{pmatrix} 33/5 \\ 1 \\ -3 \end{pmatrix} + (n_5 + 3n_{10}) \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} & b_{ia}^{(2)} &= \begin{matrix} a : & t & b & \tau \\ \begin{pmatrix} 26/5 & 14/5 & 18/5 \\ 6 & 6 & 2 \\ 4 & 4 & 0 \end{pmatrix} \end{matrix} \\ b_{ij}^{(2)} &= \begin{pmatrix} \frac{199}{25} + \frac{7n_5}{15} + \frac{23n_{10}}{5} & \frac{27}{5} + \frac{9n_5}{5} + \frac{3n_{10}}{5} & \frac{88}{5} + \frac{32n_5}{15} + \frac{48n_{10}}{5} \\ \frac{9}{5} + \frac{3n_5}{5} + \frac{n_{10}}{5} & 25 + 7n_5 + 21n_{10} & 24 + 16n_{10} \\ \frac{11}{5} + \frac{4n_5}{15} + \frac{6n_{10}}{5} & 9 + 6n_{10} & 14 + \frac{34n_5}{3} + 34n_{10} \end{pmatrix}. \end{aligned}$$

In numerical computations it is useful to employ the RGE for $\rho_i \equiv M_i/\alpha_i$, since it starts at two-loop order

$$\frac{d}{dt} \rho_i = -\frac{g_i^2}{4\pi} \left[\sum_j b_{ij}^{(2)} g_j^2 M_j + \sum_a b_{ia}^{(2)} \lambda_a^2 A_a \right]. \quad (\text{B.3})$$

⁴Only the $\widetilde{X\tilde{X}}$ diagram of fig. 1, that needs a Majorana gaugino propagator to be non-zero, requires a more detailed treatment of the charge conjugation factors.

When $F \sim M^2$ supersymmetry is *hardly* broken in the effective theory below the messenger scale: the couplings at the supergauge gaugino vertices differ (by a numerically negligible amount) from the corresponding gauge couplings. However in this case the messengers are light so that it is sufficient to employ the one loop RGE equations below the messenger scale.

Finally the gauge contribution to the 2 loop $\overline{\text{DR}}$ RGE equations for the supersymmetric messenger masses M_n and for the ‘ F -terms’ F_n is

$$\frac{d}{dt} \ln M_n = \frac{d}{dt} \ln F_n = 2C_i(X_n)g_i^2 - \frac{1}{(4\pi)^2} \left\{ g_i^2 g_j^2 2C_i(X_n) 2C_j(X_n) + g_i^4 2C_i(X_n)b_i^{(1)} \right\}.$$

References

- [1] L. Alvarez-Gaume, M. Claudson and M.B. Wise, *Nucl. Phys.* **B207** (1982) 96;
M. Dine and W. Fishler, *Nucl. Phys.* **B204** (1982) 346.
- [2] G. Dvali, G.F. Giudice and A. Pomarol, *Nucl. Phys.* **B478** (1996) 31;
H. Murayama, *hep-ph/9705271*; M.A. Luty, *hep-ph/9706554*.
- [3] S. Dimopoulos, G. Dvali, G. Giudice and R. Rattazzi, *hep-ph/9705307*.
- [4] R. Barbieri, S. Ferrara and C.A. Savoy, *Phys. Lett.* **119B** (1982) 343;
P. Nath, R. Arnowitt and A. Chamseddine, *Phys. Rev. Lett.* **49** (1982) 970.
- [5] W. Siegel, *Phys. Lett.* **84B** (1979) 193;
D.M. Capper, D.R.T. Jones and P. van Nieuwenhuizen, *Nucl. Phys.* **B167** (1980) 479.
- [6] S.P. Martin and M.T. Vaughn, *Phys. Lett.* **318B** (1993) 331; Y. Yamada, *Phys. Rev. Lett.* **72** (1994) 25.
- [7] J. Hisano, H. Murayama and T. Goto, *Phys. Rev.* **D49** (1994) 1446.
- [8] See for example D.M. Pierce et al., *Nucl. Phys.* **B491** (1997) 3 and references therein.
- [9] M. Shifman, *Int. J. Mod. Phys.* **A11** (1996) 5761.
- [10] J. Hisano and M. Shifman, *hep-ph/9705417*; I. Jack and D.R.T. Jones, *hep-ph/9709364*.
- [11] S.P. Martin, *Phys. Rev.* **D55** (1997) 3177;
S. Dimopoulos, G. Giudice, A. Pomarol, *Phys. Lett.* **B389** (1996) 37.
- [12] see V.A. Smirnov, *Renormalization and Asymptotic Expansions*, Birkhäuser, Basel, 1991, and references therein.
- [13] P. Ciafaloni et al., *hep-ph/9710312*.
- [14] P. Ciafaloni and A. Strumia, *Nucl. Phys.* **B494** (1997) 41.
- [15] L. Hall and U. Sarid, *Phys. Rev. Lett.* **70** (1993) 2673.
- [16] G. Giudice, N.A. Hamed, M.A. Luty and R. Rattazzi, *hep-ph/9702???* to appear.
- [17] G. Giudice and R. Rattazzi, *hep-ph/9706540*.
- [18] S. Wolfram, *The Mathematica book*, 3rd ed. (Wolfram Media/Cambridge University Press, 1996).